

Study details

Research institution: Research and Extension Center, Mississippi State University

Date: 2022

Location: Mississippi, United States

Soil: Sharkey clay; pH (CaCl₂): 8.12; 1.0 ppm of B

Fertilizers: *Granubor*[®] Crop variety: AG43X0

Trial design: Randomized complete block with four repetitions. Treatments consisted of different B rates and sources. Pre-plant soil sample 0-6" depth composited by rep fresh root weights of six plants per plot at v4 growth state


Results

The treatment with *Granubor* (1 lb/acre of B) produced 10.3% more compared to the control (without B).





Average soybean grain yield (bushels per acre)







Study details

Research institution: Research and Extension Center, Mississippi State University

Date: 2022

Location: Mississippi United States

Soil: Dundee silt loam; pH (CaCl₂): 6.51; 0.5 ppm of B

Fertilizers: *Granubor*[®] Crop variety: AG46X0

Trial design: Randomized complete block with four repetitions. Treatments consisted of different B rates and sources. Pre-plant soil sample 0-6" depth composited by rep fresh root weights of six plants per plot at v4 growth state


Results

The treatment with *Granubor* (1 lb/acre of B) produced 4.3% more compared to the control (without B).





Average soybean grain yield (bushels per acre)





■ Control (without B)

■ Granubor 1 lb of B per acre





Conclusions of the *Granubor* trials

The 2022 soybean crop suffered through drought conditions.

These field studies emphasizes the importance of boron in terms of better water use efficiency and drought tolerance.



Study details

Research institution: Delta Research & Extension Center, Mississippi State University

Date: 2022

Location: Mississippi, United States

Crop variety: AG46X0 Soil: Dundee silt loam

• Pre-plant soil sample 0-6" depth composited by rep fresh root weights of six plants per plot at v4 growth state

• Soil with 0.5 ppm of B


Soil pH (CaCl₂): 6.51

• Organic matter: 0.5%

Fertilizer: Solubor, Solubor Flow, Solubor Flow +K

Trial design: Randomized complete block with four repetitions





Average soybean grain yield: Bushels per acre

Response of soybean to the application of B (applied via foliar)



- Solubor R1&R3: 0.25 lb ai/a at R1 & R3 growth stages
- Solubor Flow R1: 0.5 lbs ai/a at R1 growth stage
- Solubor Flow +K V2: 0.5 lbs ai/a at V2 growth stage
- Solubor Flow +K V2&R1: 0.25 lb ai/a at V2 & R1 growth stages
- Solubor V2: 0.5 lbs ai/a at V2 growth stage
- Solubor Flow +K R1: 0.5 lbs ai/a at R1 growth stage
- Solubor V2&R1: 0.25 lb ai/a at V2 & R1 growth stages
- Solubor Flow V2&R3: 0.25 lb ai/a at V2 & R3 growth stages
- Solubor Flow V2&R1: 0.25 lb ai/a at V2 & R1 growth stages
- Solubor V2&R3: 0.25 lb ai/a at V2 & R3 growth stages

- Solubor Flow +K R1&R3: 0.25 lb ai/a at R1 & R3 growth stages
- Solubor Flow R1&R3: 0.25 lb ai/a at R1 & R3 growth stages
- Solubor Flow *K R3: 0.5 lbs ai/a at R3 growth stage
- Solubor R1: 0.5 lbs ai/a at R1 growth stage
- Solubor Flow V2: 0.5 lbs ai/a at V2 growth stage
- Solubor Flow +K V2&R3: 0.25 lb ai/a at V2 & R3 growth stages
- Solubor R3: 0.5 lbs ai/a at R3 growth stage
- Solubor Flow R3: 0.5 lbs ai/a at R3 growth stage





Average soybean grain yield: Bushels per acre



Response of soybean to the application of B (applied via foliar)





Results

The highest yields occurred with the application of:

- 0.25 lb ai/a of Solubor at the R1 and R3 growth stages which produced 11.7% more compared to the control
- 0.50 lb ai/a of *Solubor* Flow at the R1 growth stages which produced **11.2% more** compared to the control
- 0.5 lbs ai/a of Solubor Flow +K at the V2 growth stage which produced 7% more compared to the control

