

Effect of doses and different sources of boron on corn yield in argillaceous soils

The micronutrient boron (B) has been used in Brazilian agriculture for a number of years. Several options for boron fertilizers exist on the market, which can generate some confusion for farmers and agronomists. Depending on the source of B, its solubility and dissolution time (release of B) may vary, affecting the recommended dosage and the handling of B in cultivation. Another concern is the dose of B that should be recommended in relation to the texture of the soil. Research shows that there is a high correlation between the content of clays (iron and aluminum oxyhydroxides) and the adsorption of B by these clays. The amount of B adsorbed by these minerals is greater, the higher the soil's pH. In other words, the higher the soil's pH and clay content, the greater the adsorption of B, and the lower the availability of this element in soil solution for the roots to be able to absorb it. So for more clayey soils, should the dose be higher than it would be for a more sandy-textured soil?

In order to aid in understanding the response of corn to some sources available on the market, some experiments were conducted by renowned researchers in Brazil.

Schaich (2021) conducted two fieldwork experiments (during the crops of 2019/20 and 2020/21) comparing sources and doses of B. Among the sources of B tested were the fertilizer *Granubor*[®] (15% B), granulated ulexite (10% B) and a technology based on potassium chloride (KCl) + two sources of B in the same granule (58% K₂O and 0.5% B). *Granubor* is a sodium tetraborate pentahydrate-based fertilizer and the sources of B in the KCl + B product are based on anhydrous sodium tetraborate (50%) and colemanite (50%). The experiments were carried out in the city of Cruz Alta, RS, with red latosol clayey-textured (Table 1). The experimental design used randomized blocks with four repetitions.

Gitti (2021) conducted a fieldwork experiment (during the crop of 2020/21) comparing sources and doses of B. Among the sources of B tested were the fertilizer *Granubor* (15% B) and granulated ulexite (10% B). The experiment was carried out at Fundacao MS, in the municipality of Maracaju, MS, in a clayey-textured dystroferic red latosol (Table 2). The experimental design used randomized blocks with five repetitions.

Table 1: Chemical and physical characteristics of the soil in the experimental areas before the experiments were set up in them. PhysioAtac, Cruz Alta, RS, 2020.

Exp.	Depth	pH	Ca	Mg	Al	Al+H	P	K	S
	cm	H ₂ O	 cmolc/dm ⁻³ mg/dm ⁻³		
Corn	0-20 cm	5.9	8.3	1.7	0	2.2	18	160	8.
Exp.	Depth	Clay	MO	V	CTC	Zn	CU	B	MN
	cm	%	g/dm ⁻³	%	cmolc/dm ⁻³	 mg/dm ⁻³		
Corn	0-20 cm	42	3.2	82.6	12.	3.3	6.1	0.5	4.3

¹ UNISC Analytical Center; Santa Cruz do Sul - RS. Extractors: P, K, Cu, Fe, Mn and Zn (Mehlich-1); S (ammonium acetate); Ca, Mg and Al (KCl 1N); OM (sodium dichromate); B (hot water); clay (densimeter method).

Table 2: Chemical and soil texture characterization of the experimental area at depths of 0-20 and 20-40 cm. MS Foundation, Maracaju/MS, Brazil, 2020.

Prof cmpH.....		MO	P	K	Ca	Mg	Al	H+Al	SB	T	V	
	CaCl ₂	H ₂ O	g/dm ⁻³	Mehlich mmol _c /dm ⁻³							%	
0-20	5.0	5.7	37.9	24.4	5.6	57.8	15.6	0.0	66.4	79.1	145.5	54.3	
20-40	5.2	5.9	24.7	2.4	2.1	55.6	12.5	0.0	48.3	70.3	118.6	59.3	
Prof cm	S	Zn	B	Cu	Mn	Fe	Relationship	K	Ca	Mg	H	Al	Clay
 mg/dm ⁻³						Ca/Mg % da CTC					%
0-20	18.8	10.2	0.4	6.2	176,9	19	3.7	3.8	39.73	10.7	45.7	0.0	50
20-40	18.3	0.8	0.2	6.8	97,3	20	4.4	1.8	46.8	10.6	40.7	0.0	50

*Soil sample collected on 05/20/2020. Farm Alegria plot area 1, Maracaju/MS, Brazil. Code FMS 0-20 cm 11074, and 20-40 cm 11075.




Effect of doses and different sources of boron on corn yield in argillaceous soils

Results

In the experiments carried out in Cruz Alta, RS, regardless of the source, the application of B boosted productivity. This average increase was 4.2; 2.8; 4.0 and 6.9 bags/ha, for the dose of 1, 2, 3 and 4 kg/ha of B, respectively, when compared to the control (Figure 1). This shows the importance of the annual application of B in the soil for productivity gains in summer corn, regardless of the boron content available in the soil analysis (see table 1). Even though the differences in yield were not statistically different, the numerical increase in yield from *Granubor* was the greatest among the compared sources for all the doses tested (Figure 1). In the clayey soil condition of the experiments ($\geq 42\%$ clay, see tables 1), the greatest boost in yield was with the dose of 4 kg/ha of B, for the source *Granubor*. This result shows how important it is to be familiar with the available B sources and the clay content of the soil, to better gauge what dose should be applied.


Figure 1: Response of corn to the application of B in increasing doses (1; 2; 3 and 4 kg/ha) using different sources available on the market. Average yield of 2019/20 and 2020/21 crops.



In the experiment carried out in Maracaju, MS, only the application of B with the source *Granubor* and with the doses of 1 and 4 kg/ha of B, boosted productivity. This increase was 6.9 and 2.1 bags/ha, for the dose of 1 and 4 kg/ha of B, respectively, when compared to the control (Figure 2). Even though the differences in yield were not statistically different, the numerical increase in yield from *Granubor* was the greatest among the compared sources for all the doses tested (Figure 2). In the clayey soil condition of the experiments ($\geq 50\%$ clay, see table 2), the greatest boost in yield was with the dose of 1 kg/ha of B, for the source *Granubor*.

The results of these experiments show that the best dose of *Granubor* is between 6.7 - 10 kg/ha (1 kg/ha of B).

Figure 2: Response of corn to the application of B in increasing doses (1; 2 and 4 kg/ha) using different sources available on the market. Crop of 2020/21 crops.



References: Gabriel Schaich, 2021. Physioatoc. || Douglas Gitti, 2021. Fundação MS.