



# Learn more about borate fertilizers

The use of the micronutrient boron (B) in agriculture is already widespread. However, there are several sources on the market that can confuse farmers and agronomists. Depending on the source of boron, its solubility may vary, affecting the recommended rate and management of boron in the field. With the intention of clearing up doubts about different products on the market, Barth & Haliski (2020) conducted a percolation study to compare the release rate of boron from various sources.

The work was conducted in Castro, PR, Brazil, at the laboratory of the Fundação ABC. The soil used had a clay texture. The experiment was conducted under two conditions: Soil without limestone application (pH 4.8) and with soil with limestone applied (pH 5.7). Among the sources tested were the refined boron fertilizer *Granubor*<sup>®</sup> (15% B), colemanite (12.5% B), and a potassium chloride (KCl)-based technology + two sources of boron in the same granule (58% K<sub>2</sub>O and 0.5% B). *Granubor* is a water-soluble sodium tetraborate pentahydrate fertilizer, while colemanite (calcium borate) is a mineral with low water solubility. The two sources of boron present in the compacted KCl+B product are based on anhydrous sodium tetraborate (50%) and colemanite (50%).

The results showed that the mineral colemanite only released 6.9% of boron after 140 days in the soil with pH 4.8 and 6.4% in the soil with pH 5.7. The compacted KCl+B product released 57.5% and 60.1% of boron after 140 days in soils with pH 4.8 and 5.7, respectively. The fertilizer *Granubor* released 95.5% and 92.9% after 140 days, respectively. These results show that the release rate of boron from *Granubor* is best matched to the absorption of boron in annual crops such as cotton.

Percentage of percolated boron under different pH conditions






# Response of cotton to fertilization with boron

Cotton is a crop that is very sensitive to boron deficiency. The recommended boron rate for cotton is between 2-3 kg B/ha/year. A field study was conducted by Fundação Chapadão on the 2018/2019 crop in the city of Chapadão do Sul – MS, Brazil in order to evaluate the response of the cotton crop to increasing rates of boron. The soil used was Red Dystrophic Latosol with boron content of 0.36 mg.dm<sup>-3</sup> in the layer from 0 to 20 cm and pH of 5.0.

The rates of boron were 0, 0.5, 1, 2, and 4 kg of B/ha applied at planting with 230 kg of MAP/ha. The experimental design used randomized blocks with five repetitions. The source of boron was *Granubor*. Other fertilizers were applied following the recommendation of the Fundação Chapadão.

The results showed an increase in cotton seed yield of 379 kg/ha with a rate of 1 kg B/ha, while the dose of 2 kg B/ha increased the yield by 428 kg/ha. These findings demonstrate a good response of the cotton crop to boron as well as the performance of *Granubor* for efficiently providing this nutrient to meet the needs of the crop.

## Response of the cotton crop to boron fertilization using *Granubor* fertilizer



*Granubor* is a refined boron fertilizer with 15% B. It is water soluble, making boron available at the time when the cotton crop really needs it.

### Sales Contacts

Adam Brodie – North America  
adam.brodie@riotinto.com

Lucy Meng – China  
lucy.meng@riotinto.com

Henri Ngho - SE Asia  
henri.ngoho@riotinto.com

Jaime Gonzalez – Latin America  
jaime.gonzalez2@riotinto.com

Punit Antal - India  
punit.antal@riotinto.com

### Agronomy Contacts

Wendall Boehlje – North America  
wendall.boehlje@riotinto.com

Abigail Jiang – China  
abigail.jiang@riotinto.com

Weng Kee Ch'ng - SE Asia  
wengkee.ch'ng@riotinto.com

Cleiton de Sequeira - Global  
cleiton.desequeira@riotinto.com

## Photos of boron deficiency in cotton



Source: Fundação MT, 2020

## References

Gabriel Barth and Adriano Haliski, 2020. Fundação ABC.  
Claudinei Kappes, 2019. Fundação Chapadão.