

Evaluation of boron doses on soybeans in two Paraguayan soils

To aid in understanding the response of soybeans to applications of different doses of boron, Zabini (2021) conducted field experiments comparing doses of boron (B) in two different types of soils in different locations in Paraguay. The fertilizer *Granubor*® (15% B), which is a granular sodium tetraborate pentahydrate, was tested. The experiments were carried out in the locations of Santa

Rosa del Monday (Alto Paraná) and Dr. J.E. Estigarribia (Caaguazú), Tables 1 and 2. The experimental design used randomized blocks with four repetitions. The experiments received the same amounts of NPK fertilizers, that is, 107 kg/ha of MAP in the furrows and 100 kg/ha of KCl in top dressing.

Table 1

Chemical and physical characteristics of the soil in the areas used for experiments before the implementation of the experiments

pH water	Clay	OM	Ca	Mg	K	Al	H+Al	t	T
	%		cmol _c /dm ⁻³						
5.04	42.23	2.63	4.23	1.9	0.12	0.53	5.18	6.37	11.02
V	m	P	P rem	S	B	Cu	Fe	Mn	Zn
	%		mg/dm ⁻³	mg/L	mg/dm ⁻³				
52.99	8.32	28.12	37.30	6.40	0.70	4.05	35.05	136.68	3.69

Santa Rosa del Monday (Alto Paraná) (crop 2020-2021)

Table 2

Chemical and physical characteristics of the soil in the areas used for experiments before the implementation of the experiments

pH water	Clay	OM	Ca	Mg	K	Al	H+Al	t	T
	%		cmol _c /dm ⁻³						
5.30	12.00	1.13	1.49	0.36	0.15	0.11	3.17	2.11	5.17
V	m	P	P rem	S	B	Cu	Fe	Mn	Zn
	%		mg/dm ⁻³	mg/L	mg/dm ⁻³				
38.68	5.21	35.52	46.38	3.69	0.25	1.62	87.37	88.86	2.63

Dr. J.E. Estigarribia (Caaguazú) (crop 2020-2021)

TECHNICAL BROCHURE: SOYBEANS IN PARAGUAYAN SOILS

Results

At the Dr. J.E. Estigarribia (Caaguazu) site, where the soil texture is sandy, there was an increase in productivity of 532 kg/ha (8.87 bags/ha) in the soybean crop compared to the control, with application of 14 kg/ha of *Granubor* (2 kg/ha B).

At the Dr. Santa Rosa del Monday (Alto Paraná) site, where the soil texture is clayey, there was an increase in productivity of 408 kg/ha (6.8 bags/ha) in the soybean crop compared to the control, with application of 14 kg/ha of *Granubor* (2 kg/ha B).

Figure 1

Response of soy to the application of *Granubor* in increasing doses (0, 1, 2, and 4 kg/ha). 2020/21 crop, Santa Rosa del Monday





Figure 2

Response of soy to the application of *Granubor* in increasing doses (0, 1, 2, and 4 kg/ha). 2020/21 crop, Dr. J.E. Estigarribia



Andre Zabini, 2021. Agrónomo Investigación Agrícola

About U.S. Borax

U.S. Borax is a global leader in the supply and science of borates—naturally-occurring minerals containing boron and other elements. We are 1,000 people serving our customers with more than 1,200 delivery locations globally. We supply around 30% of the world's need for refined borates from our worldclass mine in Boron, California, about 100 miles northeast of Los Angeles.

Our local agriculture experts understand the uses and benefits of boron on crops. In addition to a global sales team, we have a number of agronomists on staff to help fertilizer distributors maximize the benefits of borates in agriculture applications. Our ag team can answer individual growers' questions and concerns about their particular crop.

High quality, high reliability, high performance borate products. It's what we're known for.



agriculture.borax.com